

Drug

Take.....tablets/capsules times a day

☐ Before meal ☐ After meal

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

☐ Take ½ - 1 hr before food

☐ Take immediately after eating

☐ Take until finished

☐ May cause drowsiness

Drug

Take.....millilitre/ teaspoon / tablespoon times a day

☐ Before meal

☐ After meal

☐ Morning

☐ Noon

☐ Evening

☐ Bedtime

☐ Every hours

☐ As need

☐ Take ½ - 1 hr before food

☐ Take immediately after eating

☐ Take until finished

☐ May cause drowsiness

☐ Stable for 7 days after mixing

☐ Shake well before use

Drug

Take.....inhalation times a day

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

☐ Shake the inhaler well before each use

☐ Rinse your mouth after use the inhaler

Drug

Apply to the affected area

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

For external use only

จัดทำโดย กองเภสัชกรรม สำนักอนามัย กรุงเทพมหานคร

Drug

Instill drops into the affected eye

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

- do not share your medication with others-

- dispose of 30 days after opening -

Drug

Instill drops into the affected ear

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

- do not share your medication with others-

- dispose of 30 days after opening -

Drug

Apply to the affected eye

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

- do not share your medication with others -

- dispose of 30 days after opening -

จัดทำโดย กองเภสัชกรรม สำนักอนามัย กรุงเทพมหานคร