

Drug

Take.....tablets/capsules times a day

☐ Before meal ☐ After meal

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

☐ Take ½ - 1 hr before food

☐ Take immediately after eating

☐ Take until finished

☐ May cause drowsiness

Drug

Take.....millilitre/teaspoon/tablespoon.....times a day

☐ Before meal ☐ After meal

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

☐ Take ½ - 1 hr before food ☐ Take immediately after eating

☐ Take until finished ☐ May cause drowsiness

☐ Stable for 7 days after mixing ☐ Shake well before use

Pharmaceutical Division, Bangkok Metropolitan Administration

Drug

Take.....inhalation times a day

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

☐ Shake the inhaler well before each use

☐ Rinse your mouth after use the inhaler

Drug

Apply to the affected area

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

For external use only

Pharmaceutical Division, Bangkok Metropolitan Administration

Drug

Instill drops into the affected eye

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

-
- do not share your medication with others -
 - dispose of 30 days after opening -

Pharmaceutical Division, Bangkok Metropolitan Administration

Drug

Instill drops into the affected ear

☐ Morning ☐ Noon ☐ Evening ☐ Bedtime

☐ Every hours ☐ As need

-
- do not share your medication with others -
 - dispose of 30 days after opening -

Pharmaceutical Division, Bangkok Metropolitan Administration

Drug

Apply to the affected eye

- ☐ Morning ☐ Noon ☐ Evening ☐ Bedtime
- ☐ Every hours ☐ As need

-
- do not share your medication with others -
 - dispose of 30 days after opening -

Pharmaceutical Division, Bangkok Metropolitan Administration